

**BASE RATES OF DSM-4 AXIS I DIAGNOSES
WHITE ADOLESCENT MALES (N = 675 or 4.4% OF THE TOTAL N)**

2PT	BR-GR	Dx 1	BR 1	Dx 2	BR 2	Dx 3	BR 3	Dx 4	BR 4	Dx 5	BR 5	L	2P-HR
1-7	8.4	313.81	3.5	312.9	3.5	314.01	3.5					3	3.5%
4-1	8.3	313.81	66.1	312.9	66.1	314.01	66.1					3	66.1%
4-7	7.7	309.9	67.3	303.9	1.9	300.81	1.9	300.7	1.9	300.23	1.9	5	15.0%
1-4	5.2	313.81	74.3	312.9	74.3	314.01	74.3	300.4	2.9	296.2x	2.9	5	45.7%
4-8	5.0	312.9	58.8	314.01	58.8	295.40	58.8	297.1	8.8	296.50	8.8	5	38.8%
7-4	4.4	309.9	73.3									1	73.3%
7-8	3.7	295.40	56.0	295.60	52.0	313.0	52.0	300.02	8.0	296.2x	4.0	5	34.4%
1-2	3.4	313.0	56.5	300.4	56.5	315.9	56.5	300.81	13.0	296.2x	8.7	5	38.2%
7-1	3.4	295.40	4.3	298.8	4.3	313.x	4.3					3	4.3%
2-7	3.3	313.x	40.9	300.4	36.4	296.2x	36.4	300.02	40.9	296.50	31.8	5	37.3%
1-8	3.1	296.2x	38.1									1	38.1%
8-7	3.0	295.40	25.0	295.6x	25.0	313.x	25.0	300.02	5.0	296.2x	5.0	5	17.0%
7-2	2.8	313.x	52.6	300.4	52.6	296.2x	47.4	300.02	52.6			4	51.3%
4-5	2.8	309.3	52.6	300.81	5.3	309.4	21.0					3	26.3%
8-2	2.8	296.50	78.9	296.2x	78.9	296.3x	73.7	300.4	73.4	295.7	73.7	5	75.7%
1-3	2.5	300.81	5.9	313.x	5.9							2	5.9%
2-4	2.2	309.0	40.0	300.4	40.0	305.9	40.0	296.2x	40.0	296.3x	40.0	5	40.0%
4-2	2.1	309.0	42.9	305.0	35.7	305.9	42.9	296.2x	42.9	296.3x	35.7	5	40.0%
9-4	2.1	312.9x	50.0	312.9x	50.0	309.0	57.1	309.3	7.1			4	41.0%
8-4	1.9	305.9	61.5	295.3x	53.8	295.90	46.1	297.1	53.8	296.50	53.8	5	53.8%
4-9	1.3	312.9x	33.3	314.01	11.1	309.0	33.3					3	33.3%
1-5	1.2												
2-8	1.2												
5-7	1.2												
2-1	1.0												
8-1	1.0												
8-6	1.0												
RDx-HR			46.7%		37.5%		41.1%		26.9%		24.2%	4	37.1%

- 2PT** = The most frequent 2-point patterns for the population grp. **BR 1-5** = Base rate of the most frequent DSM-4 Axis I Dx
BR = Base rates of the 2-point patterns **2P-HR** = Individual 2-pt Pattern Hit-Rates
Dx 1-5 = Ranking diagnoses from total number of episodes of care **RDx-HR** = Average Base Rate per Ranking DSM-IV Dx